VOL. 515 JOURNAL OF CHROMATOGRAPHY AUGUST 31, 1990

CONTENTS

INTERNATIONAL SYMPOSIUM ON CHROMATOGRAPHY (CIS'89), TOKYO, OCTOBER 17–20, 1989

Foreword
by T. Okuyama (Tokyo, Japan), S. Honda (Higashi-Osaka, Japan) and T. Hoshino and H. Nakamura (Tokyo, Japan) 1

Solubility parameter treatment for the characterization of the stationary phase in the reversed-phase chromatography of benzene derivatives
by F. M. Yamamoto and S. Rokushika (Kyoto, Japan) 3

Advances in expert systems for high-performance liquid chromatography
by Y. Zhang, H. Zou and P. Lu (Dalian, China) 13

Effects of ionic strength of eluent on size analysis of sub-micrometre particles by sedimentation field-flow fractionation
by Y. Mori (Kyoto, Japan) and B. Scarlett and H. G. Merkus (Delft, The Netherlands) . 27

Size-exclusion chromatography dimension for rod-like macromolecules
by P. L. Dubin, J. I. Kaplan, B.-S. Tian and M. Mehta (Indianapolis, IN, U.S.A.) . . . 37

Preparation and evaluation of octadecyl-treated porous glasses. Application to the determination of methotrexate in serum
by M. Okamoto, I. Yoshida and M. Utsumi (Gifu, Japan), K. Nobuhara (Aichi, Japan) and K. Jinno (Toyohashi, Japan) 43

Novel silica-based strong anion exchanger for single-column ion chromatography
by C.-E. Lin, Y.-H. Yang and M.-H. Yang (Taipei, Taiwan) 49

Characterization of an internal-surface reversed-phase silica support for liquid chromatography and its application to assays of drugs in serum
by J. Haginaka, J. Wakai, N. Yasuda, H. Yasuda and Y. Kimura (Nishinomiya, Japan) . 59

Equilibrium of octadecylsilica gel with sodium dodecyl sulphate
by S. Hori, K. Ohtani-Senuma, S. Ohtani, K. Miyasaka and T. Ishikawa (Tokyo, Japan) . 67

Method for the preparation of internal-surface reversed-phase packing materials starting from alkylsilylated silica gels
by K. Kimata (Kyoto, Japan), R. Tsuboi (Mukoh, Japan) and K. Hosoya, N. Tanaka and T. Araki (Kyoto, Japan) 73

Optical resolution of racemic compounds on chiral stationary phases of modified cellulose
by Y. Fukui, A. Ichida, T. Shibata and K. Mori (Himeji, Japan) 85

Fundamental study of hydroxyapatite high-performance liquid chromatography. II. Experimental analysis on the basis of the general theory of gradient chromatography
by T. Kawasaki, M. Niikura and Y. Kobayashi (Tokyo, Japan) 91

Fundamental study of hydroxyapatite high-performance liquid chromatography. III. Direct experimental confirmation of the existence of two types of adsorbing surface on the hydroxyapatite crystal
by T. Kawasaki, M. Niikura and Y. Kobayashi (Tokyo, Japan) 125

Titania and zirconia: possible new ceramic microparticulates for high-performance liquid chromatography
by M. Kawahara, H. Nakamura and T. Nakajima (Tokyo, Japan) 149

Effect of pore size on the surface excess isotherm of silica packings
by K. Tani and Y. Suzuki (Kofu, Japan) 159

Copper(II)–iminodiacetic acid chelating resin as a stationary phase in the immobilized metal ion affinity chromatography of some aromatic amines
by Y. Liu and S. Yu (Beijing, China) 169

Characteristics of C_4- and C_8-bonded vinyl alcohol copolymer gels for reversed-phase high-performance liquid chromatography
by T. Ohtani, Y. Tamura, M. Kasai, T. Uchida, Y. Yanagihara and K. Noguchi (Kanagawa-ken, Japan) 175

Separation of anionic and cationic compounds of biomedical interest by high-performance liquid chromatography on porous graphitic carbon
by G. Gu and C. K. Lim (Harrow, U.K.) 183

Binding capacities of hydroxyapatite for globular proteins
by S. Inoue and N. Ohtaki (Saitama, Japan) 193

Porous glass sheets for use in thin-layer chromatography
by M. Yoshioka, H. Araki and M. Kobayashi (Osaka, Japan), F. Kaneuchi, M. Seki, T. Miyazaki, T. Utsuki, T. Yaginuma and M. Nakano (Tokyo, Japan) 205

Constant-potential amperometric detector for carbohydrates at a nickel(III) oxide electrode for micro-scale flow-injection analysis and high-performance liquid chromatography
by M. Goto (Gifu, Japan) and H. Miyahara and D. Ishii (Nagoya, Japan) 213

Enantiomeric resolution by micellar electrokinetic chromatography with chiral surfactants
by K. Otsuka (Osaka, Japan) and S. Terabe (Kyoto, Japan) 221

Ferroceneboronic acid as a derivatization reagent for the determination of brassinosteroids by high-performance liquid chromatography with electrochemical detection
by K. Gamoh and H. Sawamoto (Kochi-shi, Japan), S. Kakatsuto (Niigata, Japan) and Y. Watabe and H. Arimoto (Kyoto, Japan) 227

Chiral separation of diltiazem, trimetoquinol and related compounds by micellar electrokinetic chromatography with bile salts
by H. Nishi, T. Fukuyama and M. Matsuo (Osaka, Japan) and S. Terabe (Himeji, Japan) 233

Separation and determination of aspoxicillin in human plasma by micellar electrokinetic chromatography with direct sample injection
by H. Nishi, T. Fukuyama and M. Matsuo (Osaka, Japan) 245

High-performance liquid chromatography of thiols with differential pulse polarographic detection of the catalytic hydrogen evolution current
by X.-X. Qian, K. Nagashima, T. Hobo, Y.-Y. Guo and C. Yamaguchi (Tokyo, Japan) . 257

Detection and identification modes for the highly sensitive and simultaneous determination of various biogenic amines by coulometric high-performance liquid chromatography
by H. Takeda (Tokyo, Japan), T. Matsumiya (Kanagawa, Japan) and T. Shibuya (Tokyo, Japan) 265

Evaluation of ammonium acetate as a volatile buffer for high-performance hydrophobic-interaction chromatography
by T. Konishi, M. Kamada and H. Nakamura (Tokyo, Japan) 279

Sample introduction and elution method for preparative supercritical fluid chromatography
by Y. Yamauchi, M. Kuwajima and M. Saito (Tokyo, Japan) 285

Enrichment of eicosapentaenoic acid and docosahexaenoic acid esters from esterified fish oil by programmed extraction–elution with supercritical carbon dioxide
by S. Higashidate, Y. Yamauchi and M. Saito (Tokyo, Japan) 295

Phenols as internal standards in reversed-phase high-performance liquid chromatography in pharmaceutical analysis
by S. Yamauchi and H. Mori (Tokushima, Japan) 305

Automated high-resolution two-dimensional liquid chromatographic system for the rapid and sensitive separation of complex peptide mixtures
by K. Matsuoka, M. Taoka, T. Isobe and T. Okuyama (Tokyo, Japan) and Y. Kato (Yamaguchi, Japan) 313

Direct sample injection into the high-performance liquid chromatographic column in theophylline monitoring
by Y. Kouno and C. Ishikura (Kanagawa, Japan) and N. Takahashi, M. Homma and K. Oka (Tokyo, Japan) 321

High-performance liquid chromatography with a 3α-hydroxysteroid dehydrogenase postcolumn reactor and isoluminol–microperoxidase chemiluminescence detection
by M. Maeda, S. Shimada and A. Tsuji (Tokyo, Japan) 329

Polymeric dimethylaminopyridinium reagents for derivatization of weak nucleophiles in high-performance liquid chromatography–ultraviolet/fluorescence detection
by C. X. Gao and I. S. Krull (Boston, MA, U.S.A.) 337

Use of a focusing cylindrical lens for increasing sensitivity in the optical detector of a capillary flow-through cell
by T. Tsuda and Y. Kobayashi (Nagoya, Japan) 357

Direct injection of blood samples into a high-performance liquid chromatographic adenine analyser to measure adenine, adenosine and the adenine nucleotides with fluorescence detection
by H. Fujimori (Osaka, Japan), T. Sasaki, K. Hibi and M. Senda (Tokyo, Japan) and M. Yoshioka (Osaka, Japan) 363

Purification of D_2 dopamine receptor by photoaffinity labelling, high-performance liquid chromatography and preparative sodium dodecyl sulphate polyacrylamide gel electrophoresis
by H. Usui, Y. Takahashi, N. Maeda and H. Mitui (Niigata, Japan), T. Isobe and T. Okuyama (Tokyo, Japan) and Y. Nishizawa and S. Hayashi (Saigata, Japan) 375

Separation of amphetamines by supercritical fluid chromatography
by J.-L. Veuthey and W. Haerdi (Genève, Switzerland) 385

Determination of malonaldehyde in oxidized biological materials by high-performance liquid chromatography
by M. Tomita and T. Okuyama (Okayama, Japan) and S. Kawai (Gifu, Japan) 391

Separation of membrane protein–sodium dodecyl sulphate complexes by high-performance liquid chromatography on hydroxyapatite
by T. Hiranuma, T. Horigome and H. Sugano (Niigata, Japan) 399

Analysis of proteins by high-performance liquid chromatography with circular dichroism spectrophotometric detection
by Y. Kurosu, T. Sasaki, T. Kakakuwa, N. Sakayanagi, K. Hibi and M. Senda (Tokyo, Japan) 407

Applications of high-performance liquid chromatography in bacteriology. I. Determination of metabolites (Review)
by C. Lucarelli (Rome, Italy) and L. Radin, R. Corio and C. Eftimiadi (Genova, Italy) . 415

Determination of free catecholamines in human urine by direct injection of urine into a liquid chromatographic column-switching system with fluorimetric detection
by T. Seki (Osaka, Japan) and Y. Yanagihara and K. Noguchi (Kanagawa, Japan) . . . 435

Enantiomer separation of pyrethroid insecticides by high-performance liquid chromatography with chiral stationary phases
by N. Ôi, H. Kitahara and R. Kira (Osaka, Japan) 441

Electrochemical detection of dipeptides and dipeptide amides
by H. Tsai and S. G. Weber (Pittsburgh, PA, U.S.A.) 451

Determination of methamphetamine, amphetamine and piperidine in human urine by high-performance liquid chromatography with chemiluminescence detection
by K. Hayakawa, N. Imaizumi, H. Ishikura, E. Minogawa, N. Takayama, H. Kobayashi and M. Miyazaki (Kanazawa, Japan) 459

Rapid determination by high-performance liquid chromatography of free fatty acids released from rat platelets after derivatization with monodansylcadaverine
by Y. M. Lee, H. Nakamura and T. Nakajima (Tokyo, Japan) 467

Separation of peptide diastereomers by reversed-phase high-performance liquid chromatography and its applications. IV. New derivatization reagent for the enantiomeric analysis of α- and β-amino acids
by T. Yamada, S. Nonomura, H. Fujiwara, T. Miyazawa and S. Kuwata (Kobe, Japan) . 475

Isolation and characterization of recombinant eel growth hormone expressed in *Escherichia coli*
by S. Sugimoto, K. Yamaguchi and Y. Yokoo (Tokyo, Japan) 483

Determination of pseudouridine in human urine and serum by high-performance liquid chromatography with post-column fluorescence derivatization
by Y. Umegae, H. Nohta and Y. Ohkura (Fukuoka, Japan) 495

Identification of a new minor iridoid glycoside in *Symplocos glauca* by thermospray liquid chromatography–mass spectrometry
by J. Iida, M. Hayashi, T. Murata, M. Ono, K. Inoue and T. Fujita (Kyoto, Japan) . . . 503

Supercritical fluid extraction and chromatography of cholesterol in food samples
by C. P. Ong, H. K. Lee and S. F. Y. Li (Singapore, Singapore) 509

Supercritical fluid extraction and chromatography of steroids with Freon-22
by S. F. Y. Li, C. P. Ong, M. L. Lee and H. K. Lee (Singapore, Singapore) 515

High-performance liquid chromatography of proteins on a ceramic hydroxyapatite with volatile buffers
by T. Kadoya (Gunma, Japan) 521

o-Phthalaldehyde post-column derivatization for the determination of gizzerosine in fish meal by high-performance liquid chromatography
by H. Murakita and T. Gotoh (Tokyo, Japan) 527

Application of high-performance liquid chromatography in establishing an accurate index of blood glucose control
by T. Hoshino, Y. Takahashi and M. Suzuki (Tokyo, Japan) 531

Isolation and identification of urinary nucleosides. Applications of high-performance liquid chromatographic methods to the synthesis of 5′-deoxyxanthosine and the simultaneous determination of 5,6-dihydrouridine and pseudouridine
by K. Nakano and T. Yasaka (Osaka, Japan), K. H. Schram, M. L. J. Reimer, T. D. McClure (Tuscon, AZ, U.S.A.) and T. Nakao and H. Yamamoto (Osaka, Japan) 537

Simple method for determination of the cephalosporin DQ-2556 in biological fluids by high-performance liquid chromatography
by K. Matsubayashi, M. Yoshioka and H. Tachizawa (Tokyo, Japan) 547

Application of supercritical fluid chromatography and supercritical fluid extraction to the measurement of hydroperoxides in foods
by K. Sugiyama, T. Shiokawa and T. Moriya (Yokohama, Japan) 555

High-performance liquid chromatography of transfer ribonucleic acids on spherical hydroxyapatite beads. II. Effects of pH and sodium chloride on chromatography
by Y. Yamakawa, K. Miyasaka and T. Ishikawa (Tokyo, Japan), Y. Yamada (Tochigi, Japan) and T. Okuyama (Tokyo, Japan) 563

Isolation of the insect metabolite trehalose by high-performance liquid chromatography (Note)
by K. Iida and M. Kajiwara (Tokyo, Japan) 573

Sensitive assay system for bile acids and steroids having hydroxyl groups utilizing high-performance liquid chromatography with peroxyoxalate chemiluminescence detection
by S. Higashidate, K. Hibi, M. Senda, S. Kanda and K. Imai (Tokyo, Japan) 577

Determination of α-tocopherol, free cholesterol, esterified cholesterols and triacylglycerols in human lipoproteins by high-performance liquid chromatography
by K. Seta (Tokyo, Japan), H. Nakamura (Saitama, Japan) and T. Okuyama (Tokyo, Japan) 585

Direct determination of the antihypertensive agent Cromakalim and its major metabolites in human urine by high-performance liquid chromatography
by S. Kudoh and H. Nakamura (Tokyo, Japan) 597

Formazan derivatives as the precolumn derivatization reagents in a coupled high-performance liquid chromatographic–spectrophotometric system for trace metal determination
by H. Hoshino, K. Nakano and T. Yotsuyanagi (Sendai, Japan) 603

Separation of high-molecular mass RNAs by high-performance liquid chromatography on hydroxyapatite
by S. Hori, S. Ohtani, K. Miyasaka, T. Ishikawa and H. Tanabe (Tokyo, Japan) 611

Determination of urinary free noradrenaline by reversed-phase high-performance liquid chromatography with on-line extraction and fluorescence derivatization
by O. Nozaki and Y. Ohba (Osaka, Japan) 621

Applications of semi-micro supercritical fluid chromatography with gradient elution to synthetic oligomer separation
by M. Takeuchi and T. Saito (Tokyo, Japan) 629

Indirect fluorescence detection of sugars separated by capillary zone electrophoresis with visible laser excitation
by T. W. Garner and E. S. Yeung (Ames, IA, U.S.A.) 639

Electrochromatography with continuous sample introduction
by T. Tsuda and Y. Muramatsu (Nagoya, Japan) 645

Analysis of the components of *Paeonia radix* by capillary zone electrophoresis
by S. Honda, K. Suzuki, M. Kataoka, A. Makino and K. Kakehi (Higashi-osaka, Japan) 653

Apparatus for coupled high-performance liquid chromatography and capillary electrophoresis in the analysis of complex protein mixtures
by H. Yamamoto, T. Manabe and T. Okuyama (Tokyo, Japan) 659

Effect of polymer ion concentrations on migration velocities in ion-exchange electrokinetic chromatography
by S. Terabe and T. Isemura (Kyoto, Japan) 667

Author Index 677